


NEMO-NET - THE FLUID LENSING NEURAL NETWORK FOR GLOBAL CORAL REEF ASSESSMENT



ESTF 2020 - VED CHIRAYATH, ALAN LI, MICHAL SEGAL-ROZENHAINMER, JARRETT VAN DEN BERGH, JUAN TORRES-PEREZ, SAM PURKIS, SYLVIA EARLE NASA SILICON VALLEY, AMES RESEARCH CENTER







FLUID LENSING CAMERA






2D Fluid Lensing Algorithm, Coral Test Target, Depth = 4.5m, MSL




Flat Fluid




Raw Distorted Frames



Mean Image





2D Fluid Lensing Results




2D Fluid Lensing Integration









NEMO-NET OBJECTIVES


- I) Develop an accurate algorithm for mapping coral reefs with remote sensing at different scales.
- 2) Globally assess the present and past dynamics of coral reef systems through a large-scale active learning convolutional neural network.
- 3) Implement domain transfer learning for spectral and spatial resolution transfer learning (super resolution) across multiple sensors.
- 4) Create 3D active learning CNN training game for data training from multiple sensors.



NEMO-NET DATA SOURCES

Effective Spatial Resolution	3D	Spectral Bands	Locations
0.1 - 5 cm	YES	3	Australia, Great Barrier Reef, Pacific
0.1 - 1 cm	YES	3-7	American Samoa, Guam, Western Australia, Puerto Rico, Indo-Pacfifc
0.65 m	NO	4	US Territories
0.5 - 3 m	NO	8	Global
30 m	NO		Global
	0.1 - 1 cm 0.65 m 0.5 - 3 m	0.1 - I cm YES 0.65 m NO 0.5 - 3 m NO	0.1 - I cm YES 3-7 0.65 m NO 4 0.5 - 3 m NO 8












NEMO-NET CLASSIFICATION HIERARCHY



Level 1	Level 2	Level 3	Level 4
Zone	Biological Cover	Biological Morphology	Biological Family
Coral/Algae	Live Coral	Branching Coral	Acroporidae
Bare Substratum	Algae	Massive Coral	Poritidae
Seagrass	Seagrass	Soft Coral	Gorgoniidae
Deep water	Mangrove	Macroalgae	Merulinidae
Clouds	Bare Substratum	Turf algae	Montastraeidae
Breaking Waves	Invertebrate	Seagrass	Mussidae
Beach (Sand, Rock, Dark)	Unknown	Mangrove	Plexauridae
Terrestrial Vegetation		Bare Substratum	Agariciidae
Unknown		Invertebrate	Siderastreidae
		Unknown	Pocilloporidae
			Pectiniidae
			Fungiidae
			Green Algae
			Brown Algae
			Red Algae
			Seagrass
			Mangrove
			Bare Substratum
			Invertebrate
			Unknown

NEMO-NET FCNN ARCHITECTURE

FluidCam,
MiDAR,
WorldView,
Sentinel...



CNN structure of NeMO-Net. The encoder blocks downsamples the data as it extracts deeper and deeper features, the bridge blocks pass the intermediate feature maps to the decoder, and the decoder blocks reconstructs the classification map from the



CLASSIFICATION IS HARD & CNNS NEED TRAINING DATA



Chihuahua vs. Blueberry Muffin

Bagel vs. Sleeping Dog

Mop vs. Hungarian Sheepdog


MM-Scale Airborne
Fluid Lensing


MM-Scale Airborne
& Satellite Data


VR & App-based Active Learning & Interactive Training through IUCN, Mission Blue, & Partners

















NEMO-NET GAME








CLASSIFICATION CONVERGENCE





Heat Map of Algae






Heat Map of Coral






0% of Users Classified as Algae / Coral





PERFORMANCE METRICS: WV-2





COMPARISON TO OTHER CNNS


RGB Image



VGG16-FCN




SharpMask




Expert Classified



DeepLab v2



NeMO-Net RefineNet



Coral	Seagrass	Clouds
Sediment	Terrestrial vegetation	Wave breaking
Beach	Deep water	Other or Unknown

Method	Accuracy	Mean Precision	Mean Recall	Frequency- weighted IoU
All Classes				
VGG16-FCN	79.0%	67.4%	70.0%	66.0%
DeepLab	73.4%	55.2%	54.7%	59.4%
SharpMask	73.2%	57.4%	54.1%	60.2%
NeMO-Net (RefineNet)	83.3%	64.9%	65.8%	71.5%
Coral, sediment, and seagrass only				
VGG16-FCN	82.8%	80.1%	85.4%	70.7%
DeepLab	77.4%	68.5%	82.2%	64.1%
SharpMask	81.6%	77.9%	81.9%	69.1%
NeMO-Net (RefineNet)	85.1%	86.8%	87.9%	74.0%
KSLOF Ecognition Prediction	48.9%	56.9%	44.5%	26.2%

NEW PUBLICATIONS

- 1) Chirayath, Ved, and Alan Li. "Next-Generation Optical Sensing Technologies for Exploring Ocean Worlds-NASA FluidCam, MiDAR, and NeMO-Net." Frontiers in Marine Science 6 (2019): 521.
- 2) Segal-Rozenhaimer, Michal et al. "Cloud detection algorithm for multi-modal satellite imagery using CNNs." Remote Sensing of Environment 237 (2020).
- 3) "Drone takes to the skies to image offshore reefs." Nature (2019) Vol 570. 545.
- 4) Chirayath, Ved and Instrella, Ron. "Fluid lensing and machine learning for centimeter-resolution airborne assessment of coral reefs in American Samoa." Remote Sensing of Environment 235 (2019): 111475.






NEMO-NET LAUNCH - EARTH DAY 2020



nemonet.infc



THANK YOU!



NEMO-NET DATA PRODUCTS

Raw Data

Fluid distortions
Optical absorption
Meter-scale spatial resolution

